For Clinic Office Use Only		
Date:	Reviewed By:	
Epinephrine Expiration:		



Summer 2021

Allergic Reaction Care Plan

DOB:	Grade (2020-2021)		
Allergic to:	Previous Anaphylaxis: Yes □ No □		
Check symptoms student has had with	past reactions:		
Swelling of eyes, face, lips, throat, tongue	Skin flushing or extreme paleness	Rapid or weak pulse	
Red, raised rash anywhere on body	☐ Blueness around mouth	Hoarseness	
Difficulty in breathing	Absence of breathing	☐ Tightness in chest	
☐ Itching with or without hives	Abdominal pain	Nausea	
Loss of consciousness	Sense of impending disaster	Vomiting	
Shortness of breath	☐ Difficulty in swallowing	Dizziness	
Anxiety	☐ Fainting	Other:	
If above symptoms occur, the Student she	ould be treated with the following w	ithout delay:	
Treatment: Please check the appropriate bo	xes and sign below.		
Student is to carry Epipen/Twinject/Auvi-C	Q with backup in the Clinic.		
Student is not to carry Epipen/Twinject/Au	vi-Q.		
☐ Give Benadryl liquid. Check a	appropriate dose: 2.5MG 25M	MG □50MG	
Give epinephrine auto-injector (Epipen/ Tw A second dose of epinephrine may be given		t□.15mg □ 0.30mg t□Yes □ No	
Call 911, then call parents listed below.			
☐ Other:			
Parent/Guardian (call first)	Parent/Guardian (call second)	
Name	Name		
Phone			
Emergency Contacts (contacted only if un	able to reach both parents)		
Name	Name	Name	
Phone	Phone		
Parent Signature		Date	